

[First Hit](#) [Fwd Refs](#)[Previous Doc](#)[Next Doc](#)[Go to Doc#](#)

End of Result Set



Generate Collection

Print

L1: Entry 1 of 1

File: USPT

Apr 29, 2003

US-PAT-NO: 6555339

DOCUMENT-IDENTIFIER: US 6555339 B1

TITLE: Non-endogenous, constitutively activated human protein-coupled receptors

DATE-ISSUED: April 29, 2003

INVENTOR-INFORMATION:

NAME	CITY	STATE	ZIP CODE	COUNTRY
Liaw; Chen W.	San Diego	CA		
Behan; Dominic P.	San Diego	CA		
Chalmers; Derek T.	Solana Beach	CA		

US-CL-CURRENT: [435/69.1](#); [435/252.3](#), [435/320.1](#), [435/325](#), [435/455](#), [435/68.1](#), [530/300](#), [530/333](#), [530/350](#), [536/23.1](#), [536/24.31](#), [702/19](#), [702/20](#), [702/22](#)

CLAIMS:

What is claimed is:

1. A method for creating a non-endogenous, constitutively active version of an endogenous human G protein coupled receptor (GPCR), said endogenous GPCR comprising a transmembrane 6 region and an intracellular loop 3 region, the method consisting essentially of: (a) selecting an endogenous human GPCR comprising a proline residue in the transmembrane-6 region; (b) identifying the endogenous 16^{sup}.th amino acid residue from the proline residue of step (a), in a carboxy-terminus to amino-terminus direction; (c) altering only the identified amino acid residue of step (b) to a non-endogenous amino acid residue to create a non-endogenous version of the endogenous human GPCR; and (d) determining if the non-endogenous version of the endogenous human GPCR of step (c) is constitutively active by measuring a difference in an intracellular signal measured for the non-endogenous version as compared with a signal induced by the endogenous human GPCR.

2. The method of claim 1 wherein the amino acid residue that is two residues from said proline residue in the transmembrane 6 region, in a carboxy-terminus to amino-terminus direction, is tryptophan.

[Previous Doc](#)[Next Doc](#)[Go to Doc#](#)